

On the Computational Complexity of the Forcing Chromatic Number

Frank Harary* Wolfgang Slany† Oleg Verbitsky‡

30 August 2005

Abstract

We consider vertex colorings of graphs in which adjacent vertices have distinct colors. A graph is s -chromatic if it is colorable in s colors and any coloring of it uses at least s colors. The forcing chromatic number $F_\chi(G)$ of an s -chromatic graph G is the smallest number of vertices which must be colored so that, with the restriction that s colors are used, every remaining vertex has its color determined uniquely. We estimate the computational complexity of $F_\chi(G)$ relating it to the complexity class US introduced by Blass and Gurevich. We prove that recognizing if $F_\chi(G) \leq 2$ is US-hard with respect to polynomial-time many-one reductions. Moreover, this problem is coNP-hard even under the promises that $F_\chi(G) \leq 3$ and G is 3-chromatic. On the other hand, recognizing if $F_\chi(G) \leq k$, for each constant k , is reducible to a problem in US via a disjunctive truth-table reduction.

Similar results are obtained also for forcing variants of the clique and the domination numbers of a graph.

1 Introduction

The vertex set of a graph G will be denoted by $V(G)$. An s -coloring of G is a map from $V(G)$ to $\{1, 2, \dots, s\}$. A coloring c is *proper* if $c(u) \neq c(v)$ for any adjacent vertices u and v . A graph G is s -colorable if it has a proper s -coloring. The minimum s for which G is s -colorable is called the *chromatic number* of G and denoted by $\chi(G)$. If $\chi(G) = s$, then G is called s -chromatic.

A *partial coloring* of G is any map from a subset of $V(G)$ to the set of positive integers. Suppose that G is s -chromatic. Let c be a proper s -coloring and p be a partial coloring of G . We say that p *forces* c if c is a unique extension of p to a proper s -coloring. The domain of p will be called a *defining set for c* . We call

*Computer Science Department, New Mexico State University, Las Cruces, NM 88003, USA.

†Institut für Softwaretechnologie, Technische Universität Graz, A-8010 Graz, Austria.

‡Institut für Informatik, Humboldt Universität Berlin, D-10099 Berlin, Germany. Supported by an Alexander von Humboldt fellowship.

$D \subseteq V(G)$ a *forcing set* in G if this set is defining for some proper s -coloring of G . The minimum cardinality of a forcing set is called the *forcing chromatic number* of G and denoted by $F_\chi(G)$. This graph invariant was introduced by Harary in [21]. Here we study its computational complexity.

To establish the hardness of computing $F_\chi(G)$, we focus on the respective slice decision problems which are defined for each non-negative integer k as follows:

FORCE $_\chi(k)$

Given: a graph G .

Decide if: $F_\chi(G) \leq k$.

The cases of $k = 0$ and $k = 1$ are tractable. It is clear that $F_\chi(G) = 0$ iff $\chi(G) = 1$, that is, G is empty. Furthermore, $F_\chi(G) = 1$ iff $\chi(G) = 2$ and G is connected, that is, G is a connected bipartite graph. Thus, we can pay attention only to $k \geq 2$. Since there is a simple reduction of FORCE $_\chi(k)$ to FORCE $_\chi(k + 1)$ (see Lemma 3.2), it would suffice to show that even FORCE $_\chi(2)$ is computationally hard. This is indeed the case.

Let 3COL denote the set of 3-colorable graphs and U3COL the set of those graphs in 3COL having a unique, up to renaming colors, proper 3-coloring. First of all, note that a hardness result for FORCE $_\chi(2)$ is easily derivable from two simple observations:

$$\begin{aligned} &\text{If } F_\chi(G) \leq 2, \text{ then } G \in \text{3COL}; \\ &\text{If } G \in \text{U3COL}, \text{ then } F_\chi(G) \leq 2. \end{aligned} \tag{1}$$

The set 3COL was shown to be NP-complete at the early stage of the NP-completeness theory in [40, 17] by reduction from SAT, the set of satisfiable Boolean formulas. It will be benefittable to use a well-known stronger fact: There is a polynomial-time many-one reduction p from SAT to 3COL which is *parsimonious*, that is, any Boolean formula Φ has exactly as many satisfying assignments to variables as the graph $p(\Phi)$ has proper 3-colorings (colorings obtainable from one another by renaming colors are not distinguished). In particular, if Φ has a unique satisfying assignment, then $p(\Phi) \in \text{U3COL}$ and hence $F_\chi(p(\Phi)) \leq 2$, while if Φ is unsatisfiable, then $p(\Phi) \notin \text{3COL}$ and hence $F_\chi(p(\Phi)) > 2$.

Valiant and Vazirani [42] designed a polynomial-time computable randomized transformation r of the set of Boolean formulas such that, if Φ is a satisfiable formula, then with a non-negligible probability the formula $r(\Phi)$ has a unique satisfying assignment, while if Φ is unsatisfiable, then $r(\Phi)$ is surely unsatisfiable. Combining r with the parsimonious reduction p of SAT to 3COL, we arrive at the conclusion that FORCE $_\chi(2)$ is NP-hard with respect to *randomized* polynomial-time many-one reductions. As a consequence, the forcing chromatic number is not computable in polynomial time unless any problem in NP is solvable by a polynomial-time Monte Carlo algorithm with one-sided error.

We aim at determining the computational complexity of $F_\chi(G)$ more precisely. Our first result establishes the hardness of FORCE $_\chi(2)$ with respect to *deterministic* polynomial-time many-one reductions. The latter reducibility concept will be default in what follows. The complexity class US, introduced by Blass and Gurevich [4], consists of languages L for which there is a polynomial-time nondeterministic Turing




Figure 1: Location of the slice decision problems for $F_\chi(G)$ in the hierarchy of complexity classes.

machine N such that a word x belongs to L iff N on input x has exactly one accepting computation path. Denote the set of Boolean formulas with exactly one satisfying assignment by USAT. This set is complete for US. As easily seen, U3COL belongs to US and, by the parsimonious reduction from SAT to 3COL, U3COL is another US-complete set. By the Valiant-Vazirani reduction, the US-hardness under polynomial-time many-one reductions implies the NP-hardness under randomized reductions and hence the former hardness concept should be considered stronger. It is known that US includes coNP [4] and this inclusion is proper unless the polynomial time hierarchy collapses [38]. Thus, the US-hardness implies also the coNP-hardness. We prove that the problem $\text{FORCE}_\chi(2)$ is US-hard.

Note that this result is equivalent to the reducibility of U3COL to $\text{FORCE}_\chi(2)$. Such a reduction would follow from the naive hypothesis, which may be suggested by (1), that a 3-chromatic G is in U3COL iff $F_\chi(G) = 2$. It should be stressed that the latter is far from being true in view of Lemma 2.7.3 below.

On the other hand, we are able to estimate the complexity of each $\text{FORCE}_\chi(k)$ from above by putting this family of problems in a complexity class which is a natural extension of US. We show that, for each $k \geq 2$, the problem $\text{FORCE}_\chi(k)$ is reducible to a set in US via a polynomial-time disjunctive truth-table reduction (*dtt-reduction* for brevity, see Section 2 for definitions). This improves on the straightforward inclusion of $\text{FORCE}_\chi(k)$ in Σ_2^P .

Denote the class of decision problems reducible to US under dtt-reductions by $P_{\text{dtt}}^{\text{US}}$. As shown by Chang, Kadin, and Rohatgi [5], $P_{\text{dtt}}^{\text{US}}$ is strictly larger than US unless the polynomial time hierarchy collapses to its third level. The position of the problems under consideration in the hierarchy of complexity classes is shown in Figure 1, where $P^{\text{NP}[\log n]}$ denotes the class of decision problems solvable by polynomial-time Turing machines with logarithmic number of queries to an NP oracle. The latter class coincides with the class of problems polynomial-time truth-table reducible to NP, see [26]. Another relation of $P_{\text{dtt}}^{\text{US}}$ to known complexity classes is $P_{\text{dtt}}^{\text{US}} \subseteq \text{C=P} \subseteq \text{PP}$, where a language L is in PP (resp. C=P) if it is recognizable




Figure 2: The class $\{G : F_\chi(G) \leq 2\}$ and surroundings.

by a nondeterministic Turing machine M with the following acceptance criterion: an input word w is in L iff at least (resp. precisely) a half of the computing paths of M on w are accepting. This inclusion follows from the facts that $US \subseteq C=P$ and that $C=P$ is closed under dtt-reductions (see [27, Theorem 9.9]).

In a recent paper [25], Hatami and Maserrat obtain a result that, in a sense, is complementary to our work and by this reason is also shown in Figure 1. Let $FORCE_\chi(*) = \{(x, k) : F_\chi(x) \leq k\}$. The authors of [25] prove that the recognition of membership in $FORCE_\chi(*)$ is a Σ_2^P -complete problem. Note that [25] and our paper use different techniques and, moreover, the two approaches apparently cannot be used in place of one another.

Our next result gives a finer information about the hardness of $FORCE_\chi(2)$. Note that, if $\chi(G) = 2$, then $F_\chi(G)$ is equal to the number of connected components of G . It turns out that the knowledge that $\chi(G) = 3$ does not help in computing $F_\chi(G)$. Moreover, it is hard to recognize whether or not $F_\chi(G) = 2$ even if it is known that $F_\chi(G) \leq 3$. Stating these strengthenings, we relax our hardness concept from the US -hardness to the $coNP$ -hardness (as already mentioned, the former implies the latter but the converse is not true unless the polynomial time hierarchy collapses). Thus, we prove that the problem $FORCE_\chi(2)$ is $coNP$ -hard even under the promises that $G \in 3COL$ and $F_\chi(G) \leq 3$ (see Figure 2). Note that the Valiant-Vazirani reduction implies no kind of a hardness result for the promise version of $FORCE_\chi(2)$.

In fact, many other graph characteristics also have natural forcing variants. Recall that a *clique* in a graph is a set of pairwise adjacent vertices. The maximum number of vertices of G in a clique is denoted by $\omega(G)$ and called the *clique number* of G . A clique is *optimal* if it consists of $\omega(G)$ vertices. A set of vertices is called *forcing* if it is included in a unique optimal clique. We denote the minimum cardinality of a forcing set by $F_\omega(G)$ and call it the *forcing clique number* of G .

Furthermore, we say that a vertex of a graph G *dominates* itself and any adjacent vertex. A set $D \subseteq V(G)$ is called *dominating* if every vertex of G is dominated by a vertex in D . The *domination number* of G , denoted by $\gamma(G)$, is the minimum cardinality of a dominating set of G . Similarly to the above, a *forcing set* of vertices

is one included in a unique optimal dominating set. The minimum cardinality of a forcing set is denoted by $F_\gamma(G)$ and called the *forcing domination number* of G . This graph invariant is introduced and studied by Chartrand, Gavlas, Vandell, and Harary [6].

For the forcing clique and domination numbers we consider the respective slice decision problems $\text{FORCE}_\omega(k)$ and $\text{FORCE}_\gamma(k)$ and show the same relation of them to the class US that we have for the forcing chromatic number. Actually, the dtt-reducibility to US is proved for all the three numbers by a uniform argument. However, the US-hardness with respect to many-one reductions for ω and γ is proved differently than for χ . The case of ω and γ seems combinatorially simpler because of the following equivalence: A graph G has a unique optimal clique iff $F_\omega(G) = 0$ and similarly with γ . The study of *unique optimum (UO) problems* was initiated by Papadimitriou [36]. Due to the US-hardness of the UO CLIQUE and UO DOMINATING SET problems, we are able to show the US-hardness of $\text{FORCE}_\omega(k)$ and $\text{FORCE}_\gamma(k)$ using only well-known standard reductions, whereas for $\text{FORCE}_\chi(k)$ we use somewhat more elaborate reductions involving graph products.

Overview of previous related work

Forcing chromatic number of particular graphs. Let K_n (resp. C_n , P_n) denote the complete graph (resp. the cycle, the path) on n vertices. As a simple exercise, we have $F_\chi(C_{2m+1}) = m + 1$. Mahmoodian, Naserasr, and Zaker [34] compute the forcing chromatic number of several Cartesian products: $F_\chi(C_{2m+1} \times K_2) = m + 1$, $F_\chi(C_m \times K_n) = m(n - 3)$ and $F_\chi(P_m \times K_n) = m(n - 3) + 2$ if $n \geq 6$, and $F_\chi(K_m \times K_n) = m(n - m)$ if $n \geq m^2$. Mahdian, Mahmoodian, Naserasr, and Harary [33] determine a few missing values: $F_\chi(C_m \times K_3) = \lfloor m/2 \rfloor + 1$ and, if m even, $F_\chi(C_m \times K_5) = 2m$. On the other hand, the asymptotics of $F_\chi(K_n \times K_n)$ are unknown (a problem having arisen in research on Latin squares, see below). The best lower and upper bounds $\lfloor 4(n - 2)/3 \rfloor \leq F_\chi(K_n \times K_n) \leq n^2/4$ are obtained, respectively, in [28] for $n \geq 8$ and [14] for all n . Our results show that no general approach for efficient computing the forcing chromatic number is possible unless $\text{NP} = \text{P}$ (and even $\text{US} = \text{P}$).

Latin squares and the complexity of recognizing a forcing coloring. A *Latin square of order n* is an $n \times n$ matrix with entries in $\{1, 2, \dots, n\}$ such that every row and column contains all the n numbers. In a *partial Latin square* some entries may be empty and every number occurs in any row or column at most once. A partial Latin square is called a *critical set* if it can be completed to a Latin square in a unique way. Colbourn, Colbourn, and Stinson [11] proved that recognition if a given partial Latin square L is a critical set is coNP-hard. Moreover, the problem is coNP-complete even if one extension of L to a Latin square is known. The result of [11] is strengthened in [16].

As it is observed in [34], there is a natural one-to-one correspondence between Latin squares of order n and proper n -colorings of the Cartesian square $K_n \times K_n$

which matches critical sets and forcing colorings.¹ It follows that it is coNP-hard to recognize if a given partial coloring p of a graph is forcing. Moreover, even if this problem is restricted to graphs $K_n \times K_n$ and one extension of p to a proper n -coloring is given, the problem is coNP-complete (see also Proposition 2.8 below).

Complexity of the forcing matching number Given a graph G with a perfect matching, Harary, Klein, and Živković [23] define its forcing matching number as the minimum size of a forcing set of edges, where the latter is a set contained in a unique perfect matching. Denote this number by $F_\nu(G)$. Let $\text{FORCE}_\nu(k)$ be the problem of recognition, given G , if $F_\nu(G) \leq k$ and let $\text{FORCE}_\nu(*)$ be the variant of the same problem with k given as a part of an input. From the polynomial-time solvability of the perfect matching problem, it easily follows that each $\text{FORCE}_\nu(k)$ is polynomial-time solvable and that $\text{FORCE}_\nu(*)$ is in NP. Afshani, Hatami, and Mahmoodian [3], using an earlier result by Adams, Mahdian, and Mahmoodian [1], prove that the latter problem is NP-complete.

Variety of combinatorial forcing numbers. Critical sets are studied since the seventies (Nelder [35]). The forcing chromatic, domination, and matching numbers attracted attention of researchers in the nineties. In fact, a number of other problems in diverse areas of combinatorics have a similar forcing nature. Defining sets in block designs (Gray [18]) have a rather rich bibliography. Other graph invariants whose forcing versions have appeared in the literature are the orientation number (Chartrand, Harary, Schultz, and Wall [7]), the geodetic number, the hull number, the dimension of a graph (Chartrand and Zhang, respectively, [8, 9, 10]), and this list is still inexhaustive. As a general concept, the combinatorial forcing was presented by Harary in a series of talks, e.g. [21].

Unique colorability. The concept of a uniquely colorable graph was introduced by Harary, Hedetniemi, and Robinson [22]. Complexity-theoretic concepts related to this combinatorial phenomenon were introduced by Valiant and Vazirani [42] and Blass and Gurevich [4]. However, the complexity-theorists prefer to deal with USAT, the uniqueness version of the archetypical SATisfiability problem. The results established for USAT, as US- and coNP-hardness under many-one reductions and NP-hardness under randomized reductions, carry through for U3COL by means of the parsimonious many-one reduction of SAT to 3COL. A direct, purely combinatorial way to show the coNP-hardness of U3COL is given by a result of Greenwell and Lovász ([19], see also [29, Theorem 8.5]). They prove, in particular, that if a connected graph G is not 3-colorable then the categorical product $G \cdot K_3$ is a uniquely colorable graph. On the other hand, if G is 3-colorable then $G \cdot K_3$ can be colored in two different ways. The latter follows from a simple observation: any proper 3-coloring of one of the factors efficiently induces a proper 3-coloring of the product $G_1 \cdot G_2$. By the NP-completeness of 3COL, we arrive at the conclusion that

¹Such a correspondence is also well known for edge colorings of the complete bipartite graph $K_{n,n}$.

the problem of recognizing, given a graph H and its proper 3-coloring, whether or not H is uniquely 3-colorable, is coNP-complete. Later this complexity-theoretic fact was observed by Dailey [15] who uses an identical combinatorial argument.

Organization of the paper

In Section 2 we define the categorical and the Cartesian graph products, which will play an important role in our proofs, and prove some preliminary lemmas about forcing sets in product graphs. We also state a few basic bounds for $F_\chi(G)$ and determine the complexity of recognition if a given set of vertices in a graph is forcing (the latter result is not used in the sequel but is worth being noticed). The hardness of $\text{FORCE}_\chi(k)$ is established in Section 3. A closer look at $\text{FORCE}_\chi(2)$ is taken in Section 4. In Section 5, using a combinatorial result of Hajiabolhassan, Mehrabadi, Tusserkani, and Zaker [20], we analyze the complexity of a related graph invariant, namely, the largest cardinality of an inclusion-minimal forcing set. Before taking into consideration the forcing clique and domination numbers, we suggest a general setting for forcing combinatorial numbers in Section 6. It is built upon the standard formal concept of an NP optimization problem. We benefit from this formal framework in some proofs. Section 7 is devoted to the forcing clique and domination numbers. The dtt-reducibility of $\text{FORCE}_\pi(k)$ to US for $\pi \in \{\chi, \omega, \gamma\}$ is shown in Section 8. Section 9 contains a concluding discussion and some open questions.

2 Background

2.1 Basics of complexity theory

We suppose that the discrete structures under consideration are encoded by binary words. For example, graphs are naturally representible by their adjacency matrices. The set of all binary words is denoted by $\{0, 1\}^*$. A decision problem is identified with a language, i.e., a subset of $\{0, 1\}^*$, consisting of all yes-instances. A *many-one reduction* of a problem X to a problem Y is a map $r : \{0, 1\}^* \rightarrow \{0, 1\}^*$ such that $x \in X$ iff $r(x) \in Y$. If $r(x)$ is computable in time bounded by a polynomial in the length of x , the reduction is called *polynomial-time*. We write $X \leq_m^P Y$ to say that there is a polynomial-time many-one reduction from X to Y .

Let C be a class of decision problems (or languages). A problem Z is called *C-hard* if any X in C is $\leq_m^P$ -reducible to Z . A problem Z is called *C-complete* if Z is C-hard and belongs to C . USAT and U3COL are examples of US-complete problems.

A *disjunctive truth-table reduction* (or *dtt-reduction*) of a language X to a language Y is a transformation which takes any word x to a set of words $y_1, \dots, y_m$ so that $x \in X$ iff $y_i \in Y$ for at least one $i \leq m$. We write $X \leq_{\text{dtt}}^P Y$ to say that there is such a polynomial-time reduction from X to Y .

If C is a class of languages and $\leq$ is a reducibility, then $C \leq X$ means that $Y \leq X$ for all Y in C (i.e., X is C-hard under $\leq$) and $X \leq C$ means that $X \leq Y$ for some Y in C .

The formal framework of promise problems is developed in [39]. Let Y and Q be languages. Whenever referring to a *decision problem Y under the promise Q* , we mean that membership in Y is to be decided only for inputs in Q . A reduction r of an ordinary decision problem X to a problem Y under the promise Q is a usual many-one reduction from X to Y with the additional requirement that $r(x) \in Q$ for all x . This definition allows us to extend the notion of C -hardness to promise problems.

A polynomial-time computable function $h : \{0, 1\}^* \times \{0, 1\}^* \rightarrow \{0, 1\}^*$ is called an *AND₂ function for a language Z* if for any pair x, y we have both x and y in Z iff $h(x, y)$ is in Z . Such an h is an *OR₂ function for Z* if we have at least one of x and y in Z iff $h(x, y)$ is in Z .

2.2 Graph products

Let $E(G)$ denote the set of edges of a graph G . Given two graphs G_1 and G_2 , we define a product graph on the vertex set $V(G_1) \times V(G_2)$ in two ways. Vertices (u_1, u_2) and (v_1, v_2) are adjacent in the *Cartesian product* $G_1 \times G_2$ if either $u_1 = v_1$ and $\{u_2, v_2\} \in E(G_2)$ or $u_2 = v_2$ and $\{u_1, v_1\} \in E(G_1)$. They are adjacent in the *categorical product* $G_1 \cdot G_2$ if both $\{u_1, v_1\} \in E(G_1)$ and $\{u_2, v_2\} \in E(G_2)$.

A set $V(G_1) \times \{v\}$ for $v \in V(G_2)$ will be called a *G_1 -layer of v* and a set $\{u\} \times V(G_2)$ for $u \in V(G_1)$ will be called a *G_2 -layer of u* .

Lemma 2.1 (Sabidussi, see [29, Theorem 8.1]) $\chi(G \times H) = \max\{\chi(G), \chi(H)\}$.

If c is a proper coloring of G , it is easy to see that $c^*(x, y) = c(x)$ is a proper coloring of the categorical product $G \cdot H$. We will say that c *induces* c^* . Similarly, any proper coloring of H induces a proper coloring of $G \cdot H$. This implies the following well-known fact.

Lemma 2.2 $\chi(G \cdot H) \leq \min\{\chi(G), \chi(H)\}$.

The next proposition shows that the Cartesian and the categorical products are, respectively, AND₂ and OR₂ functions for 3COL (see [31] for an exposition of AND and OR functions).

Proposition 2.3 (.1) $G \times H \in 3\text{COL}$ iff both G and H are in 3COL.

(.2) $G \cdot H \in 3\text{COL}$ iff at least one of G and H is in 3COL.

Proof. Item 1 is straightforward by Lemma 2.1. However, Item 2 does not follow solely from Lemma 2.2 because the equality $\chi(G \cdot H) = \min\{\chi(G), \chi(H)\}$ is still an unproven conjecture made by Hedetniemi. Luckily, the conjecture is known to be true for 4-chromatic graphs (El-Zahar and Sauer, see [29, Section 8.2]), and this suffices for our claim. ■

2.3 Preliminary lemmas

Lemma 2.4 (Greenwell-Lovász [19]) *Let G be a connected graph with $\chi(G) > n$. Then $G \cdot K_n$ is uniquely n -colorable.*

The proof can be found also in [29, Theorem 8.5]. We will use not only Lemma 2.4 itself but also a component of its proof stated as the next lemma. We call a partial coloring *injective* if different colored vertices receive different colors.

Lemma 2.5 *Let G be a connected graph and p be an injective coloring of a K_n -layer of $G \cdot K_n$. Then p forces the K_n -induced coloring of $G \cdot K_n$.*

Proof. Suppose that p is an injective coloring of a K_n -layer of $u \in V(G)$ and that a proper n -coloring c of $G \cdot K_n$ extends p . Let $v \in V(G)$ be adjacent to u . Inferably, $c(v, i) = c(u, i)$ for any $i \in V(K_n)$. Since G is connected, c is forced by p to be monochrome on each G -layer. ■

Lemma 2.6 *Any proper 3-coloring of $K_3 \cdot K_3$ is induced by one of the two factors K_3 .*

Proof. Let $V(K_3) = \{1, 2, 3\}$ and denote $L_i = \{(i, 1), (i, 2), (i, 3)\}$. Suppose that c is a coloring of $K_3 \cdot K_3$. Consider it on one of the layers, say, L_2 . If $|c(L_2)| = 3$, then c is induced by the second factor on the account of Lemma 2.5. Assume that $|c(L_2)| = 2$, for example, $c(2, 1) = 1$, $c(2, 2) = 2$, and $c(2, 3) = 2$. Then $c(1, 2) = c(3, 3) = 3$ is forced, contradictory to the fact that these vertices are adjacent. There remains the possibility that $|c(L_2)| = 1$, for example, $c(2, 1) = c(2, 2) = c(2, 3) = 2$. Color 2 can occur neither in L_1 nor in L_3 . Assume that one of these layers has both colors 1 and 3, say, $c(1, 1) = 1$ and $c(1, 2) = 3$. This forces $c(3, 3) = 2$, a contradiction. Thus, $|c(L_i)| = 1$ is possible only if L_1 , L_2 , and L_3 are all monochrome and have pairwise distinct colors. This is the coloring induced by the first factor. ■

2.4 A few basic bounds

We call two s -colorings *equivalent* if they are obtainable from one another by permutation of colors. Proper s -colorings of a graph G are equivalent if they determine the same partition of $V(G)$ into s independent sets. Let $N_\chi(G)$ denote the number of such partitions for $s = \chi(G)$. Thus, $N_\chi(G)$ is equal to the number of inequivalent proper s -colorings of s -chromatic G , while the total number of such colorings is equal to $\chi(G)!N_\chi(G)$. A graph G is *uniquely colorable* if $N_\chi(G) = 1$. In particular, $G \in \text{U3COL}$ iff $\chi(G) = 3$ and $N_\chi(G) = 1$.

Lemma 2.7

$$(.1) \quad \chi(G) - 1 \leq F_\chi(G) \leq \log_2 N_\chi(G) + \log_2(\chi(G)!).$$

$$(.2) \quad \text{If } N_\chi(G) = 1, \text{ then } F_\chi(G) = \chi(G) - 1.$$




Figure 3: Proper 3-colorings of $K_3 \times K_2$.

- (.3) For any k , there is a 3-chromatic graph G_k on $4k+2$ vertices with $F_\chi(G_k) = 2$ and $N_\chi(G_k) = 2^{k-1} + 1$.

The lower bound in Item 1 is sharp by Item 2. The upper bound is sharp because, for example, $F_\chi(mK_2) = m$ while $N_\chi(mK_2) = 2^{m-1}$, where mK_2 denotes the graph consisting of m isolated edges. Item 3 shows that the converse of Item 2 is false. It is also worth noting that both the lower and the upper bounds in Item 1 are computationally hard, even if one is content with finding an approximate value. The NP-hardness of approximation of the chromatic number is established in [32]. A hardness result for approximate computing $\log_2 N_\chi(G)$, even for 3-chromatic graphs, is obtained in [43].

Proof. Item 2 and the lower bound in Item 1 are obvious. To prove the upper bound in Item 1, we have to show that a k -chromatic graph G has a forcing set of at most $l = \lfloor \log_2(k!N_\chi(G)) \rfloor$ vertices $v_1, v_2, \dots$. Let C_1 be the set of all $k!N_\chi(G)$ proper k -colorings of G . We choose vertices $v_1, v_2, \dots$ one by one as follows. Let $i \geq 1$. Assume that the preceding $i-1$ vertices have been chosen and that a set of colorings C_i has been defined and has at least 2 elements. We set v_i to be an arbitrary vertex such that not all colorings in C_i coincide at v_i . Furthermore, we assign v_i a color $p(v_i)$ occurring in the multiset $\{c(v_i) : c \in C_i\}$ least frequently. Finally, we define $C_{i+1} = \{c \in C_i : c(v_i) = p(v_i)\}$. Note that $|C_{i+1}| \leq |C_i|/2$. We eventually have $|C_{i+1}| = 1$ for some $i \leq l = \lfloor \log_2 |C_1| \rfloor$. For this i , denote the single coloring in C_{i+1} by c . By construction, $v_1, \dots, v_i$ is defining for c .

To prove Item 3, consider $H = K_3 \times K_2$. This graph has two inequivalent colorings c_1 and c_2 shown in Figure 3. Let $u, v, w \in V(H)$ be as in Figure 3. Note that a partial coloring $p_1(u) \neq p_1(v)$ forces c_1 or its equivalent and that $p_2(u) = p_2(v) \neq p_2(w)$ forces c_2 .

Let G_k consist of k copies of H with all u and all v identified, that is, G_k has $4k+2$ vertices. Since the set $\{u, v\}$ stays forcing in G_k , we have $F_\chi(G_k) = 2$. If u and v are assigned the same color, we are free to assign each copy of w any of the two remaining colors. It follows that $N_\chi(G_k) = 2^{k-1} + 1$. ■

2.5 Complexity of a forcing set recognition

Proposition 2.8 *The problem of recognizing, given a graph G and two vertices $u, v \in V(G)$, whether or not $\{u, v\}$ is a forcing set is US-complete.*

Proof. Suppose that $\chi(G) \geq 3$ for else the question is efficiently decidable. Set $p(u) = 1$ and $p(v) = 2$. Obviously, $\{u, v\}$ is a forcing set iff p forces a proper 3-coloring of G . Verification of the latter condition is clearly in US.

We now describe a reduction R of the US-complete problem U3COL to the problem under consideration. If an input graph G is empty, let $R(G)$ be an arbitrary graph with two vertices which are not a forcing set. Otherwise, let u and v be the lexicographically first pair of adjacent vertices in G . We set $R(G) = (G, u, v)$. It is not hard to see that $G \in \text{U3COL}$ iff $R(G)$ consists of a graph and a 2-vertex forcing set in it. ■

3 Complexity of $F_\chi(G)$: A lower bound

Theorem 3.1 *For each $k \geq 2$, the problem $\text{FORCE}_\chi(k)$ is US-hard. Moreover, this holds true even if we consider only connected graphs.*

We first observe that the family of problems $\text{FORCE}_\chi(k)$ is linearly ordered with respect to the $\leq_m^P$ -reducibility. A simple reduction showing this does not preserve connectedness of graphs. However, if we restrict ourselves to connected graphs, we are able to show that $\text{FORCE}_\chi(2)$ remains the minimum element in this order. We then prove that $\text{FORCE}_\chi(2)$ is US-hard (even for connected graphs).

Lemma 3.2 $\text{FORCE}_\chi(k) \leq_m^P \text{FORCE}_\chi(k+1)$.

Proof. Given a non-empty graph G , we add one isolated vertex to it. Denoting the result by $G + K_1$, it is enough to notice that $F_\chi(G + K_1) = F_\chi(G) + 1$. ■

Lemma 3.3 *Let $k \geq 2$. Then $\text{FORCE}_\chi(2)$ reduces to $\text{FORCE}_\chi(k)$ even if we consider the problems only for connected graphs.*

Proof. Let G be a graph on n vertices and $m \leq n$. Writing $H = G \oplus mK_2$, we mean that

- $V(H) = \{v_1, \dots, v_n\} \cup \bigcup_{i=1}^m \{a_i, b_i\}$,
- H induces on $\{v_1, \dots, v_n\}$ a graph isomorphic to G ,
- $\{v_i, a_i\}$ and $\{a_i, b_i\}$ for all $i \leq m$ are edges of H , and
- H has no other edges.

Suppose that $\chi(G) \geq 3$ and $H = G \oplus mK_2$. Let us check that $F_\chi(H) \leq 2 + m$ if $F_\chi(G) = 2$ and $F_\chi(H) \geq 3 + m$ if $F_\chi(G) \geq 3$. This will give us the following reduction of $\text{FORCE}_\chi(2)$ to $\text{FORCE}_\chi(k)$: Given G , construct an $H = G \oplus (k-2)K_2$.

Let $F_\chi(G) = 2$. Note that G must be 3-chromatic. To show that $F_\chi(H) \leq 2 + m$, we construct a forcing set of $2 + m$ vertices. We will identify G and its copy spanned by $\{v_1, \dots, v_n\}$ in H . Let v_j and v_l be two vertices such that a partial coloring




Figure 4: Proof of Lemma 3.5 (Case 1).

$p(v_j) = 1$ and $p(v_l) = 2$ forces a proper 3-coloring c of G . Color each b_i differently from $c(v_i)$. Then $\{v_j, v_l, b_i, \dots, b_m\}$ becomes a forcing set in H .

Let $F_\chi(G) \geq 3$. Clearly, any forcing set contains all b_i 's. We hence have to show that no set $S = \{b_1, \dots, b_m, x, y\} \subset V(H)$ is forcing in H . We will assume that $\chi(G) = 3$ for else the claim is easy. Suppose that $x \in \{v_j, a_j\}$ and $y \in \{v_l, a_l\}$ for some j, l and that S is augmented with a coloring p extendable to a proper 3-coloring c of H . Let c' denote the restriction of c to $\{v_1, \dots, v_n\}$. Since c' is not a unique coloring of G compatible with p (otherwise $\{v_j, v_l\}$ would be a forcing set for G), there is another such coloring c'' . Obviously, c'' and p have a common extension to a proper coloring of H . Thus, S is not a forcing set. ■

Lemma 3.4 $\text{FORCE}_\chi(2)$ is US-hard even if restricted to connected graphs.

To prove the lemma, we describe a reduction from U3COL. Note that U3COL remains US-complete when restricted to connected graphs and that our reduction will preserve connectedness. Since the class of 2-colorable graphs is tractable and can be excluded from consideration, the desired reduction is given by the following lemma.

Lemma 3.5 Suppose that $\chi(G) \geq 3$. Then $G \in \text{U3COL}$ iff $F_\chi(G \times K_3) = 2$.

Proof. *Case 1:* $G \in \text{U3COL}$. We have to show that $F_\chi(G \times K_3) = 2$.

Fix arbitrary $u, v \in V(G)$ whose colors in the proper 3-coloring of G are different, for example, u and v can be any adjacent vertices of G . Let $V(K_3) = \{1, 2, 3\}$. Assign $p(u, 1) = 1$ and $p(v, 2) = 2$ and check that p forces a proper 3-coloring of $G \times K_3$. Assume that c is a proper 3-coloring of $G \times K_3$ consistent with p . Since c on each G -layer coincides with the 3-coloring of G up to permutation of colors, we easily infer that $c(v, 1) = c(u, 2) = 3$ (see Figure 4). This implies $c(u, 3) = 2$ and $c(v, 3) = 1$. Thus, in each G -layer we have two vertices with distinct colors, which determines colors of all the other vertices. As easily seen, the coloring obtained is really proper.

Case 2: $G \in \text{3COL} \setminus \text{U3COL}$. We have to check that $F_\chi(G \times K_3) \geq 3$.

Given a partial coloring p of two vertices $a, b \in V(G \times K_3)$, we have to show that it is not forcing. The cases that $p(a) = p(b)$ or that a and b are in the same G - or K_3 -layer are easy. Without loss of generality we therefore suppose that $p(a) = 1$, $p(b) = 2$, $a = (u, 1)$, and $b = (v, 2)$, where u and v are distinct vertices of G . Define two partial colorings of G by $c_1(u) = c_1(v) = 1$ and by $c_2(u) = 1, c_2(v) = 3$.

Subcase 2.1: Both c_1 and c_2 extend to proper 3-colorings of G . Denote the extensions by e_1 and e_2 respectively. Denote the three G -layers of $G \times K_3$ by G_1, G_2, G_3 and consider e_1, e_2 on G_1 . For each $i = 1, 2$, e_i and p agree and have a common extension to a proper coloring of $G \times K_3$. Thus, p is not forcing.

Subcase 2.2: Only c_1 extends to a proper 3-coloring of G . Since G is not uniquely colorable, there must be at least two extensions, e_1 and e_2 , of c_1 to proper 3-colorings of G_1 . As in the preceding case, e_1 and e_2 each agree with p and together with p extend two distinct colorings of $G \times K_3$.

Subcase 2.3: Only c_2 extends to a proper coloring of G . This case is completely similar to Subcase 2.2.

Case 3: $G \notin 3\text{COL}$. We have $\chi(G \times K_3) \geq 4$ by Lemma 2.1 and $F_\chi(G \times K_3) \geq 3$ by Lemma 2.7.1. ■

Theorem 3.1 immediately follows from Lemmas 3.4 and 3.3.

4 Hardness of $\text{FORCE}_\chi(2)$: A closer look

Theorem 4.1 *The problem $\text{FORCE}_\chi(2)$ is coNP-hard even under the promises that $F_\chi(G) \leq 3$ and $\chi(G) \leq 3$ and even if an input graph G is given together with its proper 3-coloring.*

Let us for a while omit the promise that $F_\chi(G) \leq 3$. Then the theorem is provable by combining the Greenwell-Lovász reduction of coNP to US (Lemma 2.4) and our reduction of US to $\text{FORCE}_\chi(2)$ (Lemma 3.5). Doing so, we easily deduce the following:

- If $\chi(G) > 3$, then $G \cdot K_3$ is uniquely 3-colorable and hence $F_\chi((G \cdot K_3) \times K_3) = 2$.
- If $\chi(G) = 3$, then $G \cdot K_3$ is 3-chromatic because it contains an odd cycle (this is an easy particular case of the aforementioned Hedetniemi conjecture). Moreover, $G \cdot K_3$ has two induced 3-colorings and hence $F_\chi((G \cdot K_3) \times K_3) \geq 3$.

To obtain Theorem 4.1 (without the promise $F_\chi(G) \leq 3$), it now suffices to make the following observation.

Lemma 4.2 $\chi((G \cdot K_3) \times K_3) = 3$ for any graph G . Moreover, a proper 3-coloring is efficiently obtainable from the explicit product representation of $(G \cdot K_3) \times K_3$.

Proof. By Lemma 2.2, we have $\chi(G \cdot K_3) \leq 3$ and hence, by Lemma 2.1, $\chi((G \cdot K_3) \times K_3) = 3$. Let $V(K_3) = \{1, 2, 3\}$ and denote $V_{i,j} = \{(v, i, j) : v \in V(G)\}$. It is not hard to see that $V_{1,1} \cup V_{2,2} \cup V_{3,3}$, $V_{1,2} \cup V_{2,3} \cup V_{3,1}$, and $V_{1,3} \cup V_{2,1} \cup V_{3,2}$ is a partition of $V((G \cdot K_3) \times K_3)$ into independent sets. ■




Figure 5: $(G \cdot K_3) \times K_3$ (Proof of Lemma 4.4)

Remark 4.3 The known facts about graph factorizations [29, Chapters 4 and 5] imply a nontrivial strengthening of Lemma 4.2 under certain, rather general conditions. Namely, if G is a connected nonbipartite graph and no two vertices of G have the same neighborhood, then we do not need to assume that $H = (G \cdot K_3) \times K_3$ is explicitly represented as a product graph because the product structure is efficiently reconstructible from the isomorphism type of H . We thank Wilfried Imrich for this observation.

To obtain the full version of Theorem 4.1, we only slightly modify the reduction: Before transforming G in $(G \cdot K_3) \times K_3$, we add to G a triangle with one vertex in $V(G)$ and two vertices new. Provided $\chi(G) \geq 3$, this does not change $\chi(G)$ and hence the modified transformation is an equally good reduction. The strengthening (the promise $F_\chi(G) \leq 3$) is given by the following lemma.

Lemma 4.4 *If a graph G is connected and contains a triangle, then $F_\chi((G \cdot K_3) \times K_3) \leq 3$.*

Proof. Let v be a vertex of a triangle T in G . Consider the product $H = G \cdot K_3$ and a partial coloring $p(v, 1) = 1$, $p(v, 2) = 2$. We claim that p forces the K_3 -induced coloring of H . Obviously, the latter is an extension of p . To show that no other extension is possible, assume that c is a proper 3-coloring of H compatible with p and consider the restriction of c on $T \cdot K_3$. By Lemma 2.6, c on $T \cdot K_3$ coincides with the coloring induced by the second factor. In particular, $c(v, 3) = 3$. Our claim now follows from the connectedness of G by Lemma 2.5.

Now, let H_1, H_2, H_3 denote the three H -layers in $H \times K_3$ and, for each $i = 1, 2, 3$, let G_{i1}, G_{i2}, G_{i3} denote the three G -layers in H_i . Let $p(v, 1, 1) = 1$, $p(v, 2, 1) = 2$ be

the forcing partial coloring for H_1 as described above (see Figure 5). Thus, p forces coloring the whole G_{1j} in color j for each $j = 1, 2, 3$. Suppose that c is a proper 3-coloring of $H \times K_3$ that agrees with p . From the product structure of $H \times K_3$ we see that, for each j , in G_{2j} there cannot occur color j . Let T^2 denote the copy of $T \cdot K_3$ in H_2 . By Lemma 2.6, c on T^2 is induced by the second factor and hence $c(v, 1, 2)$, $c(v, 2, 2)$, and $c(v, 3, 2)$ are pairwise distinct. By Lemma 2.5, each of G_{21} , G_{22} , and G_{23} is monochrome and these layers receive pairwise distinct colors. We already know that $c(G_{2j}) \neq j$. Thus, when we define p in the third point by $p(x) = 3$ for an arbitrary $x \in G_{21}$, this forces $c(G_{21}) = 3$, $c(G_{22}) = 1$, and $c(G_{23}) = 2$. Since every vertex in G_{3j} is in triangle with its clones in G_{1j} and G_{2j} , c is uniquely extrapolated on H_3 . ■

The proof of Theorem 4.1 is complete.

5 Maximum size of a minimal forcing set

Another related invariant of a graph G is the largest cardinality of an inclusion-minimal forcing set in G . We will denote this number by $F_\chi^*(G)$. A complexity analysis of $F_\chi^*(G)$ is easier owing to the characterization of uniquely colorable graphs obtained in [20].

Lemma 5.1 (Hajiabolhassan, Mehrabadi, Tusserkani, and Zaker [20]) *A connected graph G is uniquely 3-colorable iff $F_\chi^*(G) = 2$.*

Theorem 5.2 *The problem of deciding, given a graph G and its proper 3-coloring, whether or not $F_\chi^*(G) \leq 2$ is coNP-complete.*

Proof. The problem is in coNP because a no-instance of it has a certificate consisting of a proper 3-coloring c of G and a 3-vertex set $A \subset V(G)$ such that no 2-element subset B of A is defining for c . The latter fact, for each B , is certified by another proper 3-coloring c_B that agrees with c on B but differs from c somewhere outside B .

The completeness is proved by reduction from the decision problem whether or not $\chi(G) > 3$. The latter problem is coNP-complete even if restricted to connected graphs with $\chi(G) \geq 3$. Let G be a such graph. Our reduction just transforms G into the categorical product $G \cdot K_3$. If $\chi(G) > 3$, then $G \cdot K_3$ is uniquely 3-colorable by Lemma 2.4 and, by Lemma 5.1, we have $F_\chi^*(G \cdot K_3) = 2$. If $\chi(G) = 3$, then $G \cdot K_3$ has at least two inequivalent proper 3-colorings, namely, those induced by the two factors. By Lemma 5.1, we have $F_\chi^*(G \cdot K_3) \geq 3$. ■

6 General setting

In fact, many other graph characteristics also have natural forcing variants. Taking those into consideration, it will be convenient to use the formal concept of an NP optimization problem (see e.g. [12]).

Let $\{0, 1\}^*$ denote the set of binary strings. The length of a string $w \in \{0, 1\}^*$ is denoted by $|w|$. We will use notation $[n] = \{1, 2, \dots, n\}$.

An *NP optimization problem* $\pi = (\text{opt}_\pi, I_\pi, \text{sol}_\pi, v_\pi)$ (where subscript π may be omitted) consists of the following components.

- $\text{opt} \in \{\max, \min\}$ is a *type* of the problem.
- $I \subseteq \{0, 1\}^*$ is the polynomial-time decidable set of *instances* of π .
- Given $x \in I$, we have $\text{sol}(x) \subset \{0, 1\}^*$, the set of *feasible solutions* of π on instance x . We suppose that all $y \in \text{sol}(x)$ have the same length that depends only on $|x|$ and is bounded by $|x|^{O(1)}$. Given x and y , it is decidable in polynomial time whether $y \in \text{sol}(x)$.
- $v : \{0, 1\}^* \times \{0, 1\}^* \rightarrow \mathbf{N}$ is a polynomial-time computable *objective function* taking on positive integer values. If $y \in \text{sol}(x)$, then $v(x, y)$ is called the *value* of y .

The problem is, given an instance x , to compute the optimum value

$$\pi(x) = \text{opt}_{y \in \text{sol}(x)} v(x, y).$$

Such a problem is called *polynomially bounded* if $v(x, y) = |x|^{O(1)}$ for all $x \in I$ and $y \in \text{sol}(x)$.

Any $y \in \text{sol}(x)$ whose value is optimum is called an *optimum solution* of π on instance x . Let $\text{optsol}(x)$ denote the set of all such y . Given an NP optimization problem π , we define

$$\text{UO}_\pi = \{x : |\text{optsol}(x)| = 1\}.$$

Example 6.1 The problem of computing the chromating number of a graph is expressible as a quadruple $\chi = (\min, I, \text{sol}, v)$ as follows. A graph G with vertex set $V(G) = \{v_1, \dots, v_n\}$ is represented by its adjacency matrix written down row after row as a binary string x of length n^2 . A feasible solution, that is a proper coloring $c : V(G) \rightarrow [n]$, is represented by a binary string $y = c(v_1) \dots c(v_n)$ of length n^2 , where a color i is encoded by string $0^{i-1}10^{n-i}$. The value $v(x, y)$ is equal to the actual number of colors occurring in y .

Example 6.2 For the problem of computing the clique number it is natural to fix the following representation. An instance graph G is encoded as above. A feasible solution, which is a subset of $V(G)$, is encoded by its characteristic binary string of length n . The problem of computing the domination number is represented in the same way.

Given a non-empty set $U \subseteq \{0, 1\}^l$, we define $\text{force}(U)$ to be the minimum cardinality of a set $S \subseteq [l]$ such that there is exactly one string in U with 1 at every position from S . With each NP optimization problem π we associate its *forcing number* F_π , an integer-valued function of instances of π defined by

$$F_\pi(x) = \text{force}(\text{optsol}(x)).$$

Let $\text{FORCE}_\pi(k) = \{x : F_\pi(x) \leq k\}$. It is easy to check that, if χ , ω , and γ are represented as in Examples 6.1 and 6.2, then F_χ , F_ω , and F_γ are precisely those graph invariants introduced in Section 1.

Note that $\text{force}(U) = 0$ iff U is a singleton. It follows that for $\pi \in \{\omega, \gamma\}$ we have

$$x \in \text{UO}_\pi \text{ iff } F_\pi(x) = 0. \quad (2)$$

This will be the starting point of our analysis of decision problems $\text{FORCE}_\omega(k)$ and $\text{FORCE}_\gamma(k)$ in the next section.

7 Hardness of $\text{FORCE}_\omega(k)$ and $\text{FORCE}_\gamma(k)$

The results stated here are based on known reducibilities between several optimization problems related to our work. Since this material is dispersed through the literature with proofs sometimes skipped, for the reader's convenience we outline some details. We first introduce some reducibility concepts for NP optimizations problems.

Let π and ϖ be NP optimization problems of the same type. Let $f : \{0, 1\}^* \rightarrow \{0, 1\}^*$ and $g : \{0, 1\}^* \times \{0, 1\}^* \rightarrow \{0, 1\}^*$ be polynomial-time computable functions such that for every $x \in I_\pi$ we have $f(x) \in I_\varpi$ and for every $y \in \text{sol}_\varpi(f(x))$ we have $g(x, y) \in \text{sol}_\pi(x)$. Such a pair (f, g) is said to be an *S-reduction* from π to ϖ if for every $x \in I_\pi$ we have

$$\text{opt}_\pi(x) = \text{opt}_\varpi(f(x))$$

and, in addition, for every $y \in \text{sol}_\varpi(f(x))$ we have

$$v_\pi(x, g(x, y)) = v_\varpi(f(x), y).$$

We call an *S-reduction* (f, g) a *parsimonious reduction* from π to ϖ if, for any $x \in I_\pi$, $g(x, \cdot)$ is a one-to-one map from $\text{sol}_\varpi(f(x))$ onto $\text{sol}_\pi(x)$. If only a weaker condition is met, namely, that $g(x, \cdot)$ is a one-to-one correspondence between the optimum solutions of ϖ on instance $f(x)$ and the optimum solutions of π on instance x , then (f, g) will be called a *weakly parsimonious reduction* from π to ϖ .

Given a Boolean formula Φ in the conjunctive normal form (CNF), let $\sigma(\Phi)$ denote the maximum number of clauses of Φ satisfiable by the same truth assignment to the variables. By σ_3 we denote the restriction of σ to 3CNF formulas (those having at most 3 literals per clause). We regard σ and σ_3 as NP optimization problems. Both problems belong to the class MAX NP introduced by Papadimitriou and Yannakakis. Crescenzi, Fiorini, and Silvestri [13], who introduced the notion of an S-reduction, proved that every problem in MAX NP is *S-reducible* to ω , the maximum clique problem. We need a somewhat stronger fact about σ_3 .

Lemma 7.1 *There is a parsimonious reduction from σ_3 to ω .*

Proof. Let ϕ be a disjunctive clause and X be the set of variables occurring in ϕ . Let $D(\phi)$ denote the set of all conjunctions that contain every variable from X or

its negation and imply ϕ . Note that ϕ is logically equivalent to the disjunction of all ψ in $D(\phi)$.

Given a 3CNF formula Φ , we construct a graph G as follows. Let $V(G)$ be the union of $D(\phi)$ over all clauses ϕ of Φ . We join ψ_1 and ψ_2 in $V(G)$ by an edge if these conjunctions are consistent, i.e., no variable occurring in ψ_1 occurs in ψ_2 with negation and vice versa. ■

Lemma 7.2 (Thierauf [41, Section 3.2.3]) *The decision problem UO_ω is US-hard.*

Proof. Denote the restrictions of SAT and USAT to 3CNF formulas by 3SAT and U3SAT respectively. Since there is a parsimonious $\leq_m^P$ -reduction from SAT to 3SAT (see e.g. [37]), U3SAT is US-complete. We now show that $\text{U3SAT} \leq_m^P \text{UO}_\omega$.

Given a 3CNF formula Φ , let G be the graph constructed from Φ by the reduction of Lemma 7.1. Let m denote the number of clauses in Φ and $H = G + 2K_{m-1}$, the disjoint union of G and two copies of K_{m-1} .

If $\Phi \in \text{U3SAT}$, then $\omega(H) = \omega(G) = m$ and $H \in \text{UO}_\omega$ because $G \in \text{UO}_\omega$.

If $\Phi \in \text{SAT} \setminus \text{U3SAT}$, then $\omega(H) = \omega(G) = m$ and $H \notin \text{UO}_\omega$ because $G \notin \text{UO}_\omega$.

If $\Phi \notin \text{SAT}$, then $\omega(H) \leq m - 1$ and $H \notin \text{UO}_\omega$ having at least two optimal cliques.

Thus, $\Phi \in \text{U3SAT}$ iff $H \in \text{UO}_\omega$. ■

Lemma 7.3 $\text{UO}_\omega \leq_m^P \text{UO}_\gamma$.

Proof. Recall that a *vertex cover* of a graph G is a set $S \subseteq V(G)$ such that every edge of G is incident to a vertex in S . The *vertex cover number* of G is defined to be the minimum cardinality of a vertex cover of G and denoted by $\tau(G)$. It is easy to see and well known that $S \subseteq V(G)$ is a clique in G iff $V(G) \setminus S$ is a vertex cover in the graph complementary to G . It follows that

$$\text{UO}_\omega \leq_m^P \text{UO}_\tau. \quad (3)$$

We now show that

$$\text{UO}_\tau \leq_m^P \text{UO}_\gamma \quad (4)$$

by regarding τ and γ as NP minimization problems and designing a weakly parsimonious reduction from τ to γ . Recall that, given a set X and a system of its subsets $\mathcal{Y} = \{Y_1, \dots, Y_n\}$, a subsystem $\{Y_{i_1}, \dots, Y_{i_k}\}$ is called a *set cover* if $X = \bigcup_{j=1}^k Y_{i_j}$. We compose two known reductions between minimization problems, Reduction A from the minimum vertex cover to the minimum set cover ([2, Theorem 10.11]) and Reduction B from the minimum set cover to the minimum domination number (an adaptation of [30, Theorem A.1]).

Reduction A. Given a graph G and its vertex v , let $I(v)$ denote the set of the edges of G incident to v . Consider the set $X = E(G)$ and the system of its subsets $\mathcal{Y} = \{I(v) : v \in V(G)\}$. Then $S \subseteq V(G)$ is an optimal vertex cover for G iff $\{I(v) : v \in S\}$ is an optimal set cover for $(X, \mathcal{Y})$.

Reduction B. Given a set $X = \{x_1, \dots, x_m\}$ and a system of sets $\mathcal{Y} = \{Y_1, \dots, Y_n\}$ such that $X = \bigcup_{j=1}^n Y_j$, we construct a graph H as follows. $V(H)$ contains each element x_i in duplicate, namely, x_i itself and its clone x'_i . There is no edge between these $2m$ vertices. Other vertices of H are indices $1, \dots, n$, with all possible $\binom{n}{2}$ edges between them. If and only if $x_i \in Y_j$, both x_i and x'_i are adjacent to j . There are no more vertices and edges.

Observe that any optimal dominating set $D \subset V(H)$ is included in $[n]$. Indeed, if D contains both x_i and x'_i , then it can be reduced by replacing these two vertices by only one vertex j such that $x_i \in Y_j$. If D contains exactly one of x_i and x'_i , say x_i , then it should contain some j such that $x_i \in Y_j$ to dominate x'_i . But then D can be reduced just by removing x_i .

It is also clear that a set $D \subseteq [n]$ is dominating in H iff $\{Y_j : j \in D\}$ is a set cover for $(X, \mathcal{Y})$. Thus, there is a one-to-one correspondence between optimal set covers for $(X, \mathcal{Y})$ and optimal dominating sets in H . ■

Thus, UO_ω and UO_γ are both US-hard.

Lemma 7.4 *Let $\pi \in \{\omega, \gamma\}$. Then $\text{FORCE}_\pi(k) \leq_m^P \text{FORCE}_\pi(k+1)$ for any $k \geq 0$.*

Proof. Given a graph G , we have to construct a graph H such that $F_\pi(G) \leq k$ iff $F_\pi(H) \leq k+1$. It suffices to ensure that

$$F_\pi(H) = F_\pi(G) + 1. \quad (5)$$

Let $\pi = \omega$. Let H be the result of adding to G two new vertices u and v and the edges $\{w, u\}$ and $\{w, v\}$ for all $w \in V(G)$. Any optimal clique in H consists of an optimal clique in G and of either u or v . Hence any forcing set in H consists of a forcing set in G and of either u or v (we use the terminology of Section 1). This implies (5).

If $\pi = \gamma$, we obtain H from G by adding a new isolated edge. ■

Putting it all together, we make the following conclusion.

Theorem 7.5 *Let $\pi \in \{\omega, \gamma\}$. Then*

$$\text{US} \leq_m^P \text{UO}_\pi = \text{FORCE}_\pi(0) \leq_m^P \text{FORCE}_\pi(k) \leq_m^P \text{FORCE}_\pi(k+1)$$

for any $k \geq 0$.

8 Complexity of $\text{FORCE}_\pi(k)$: An upper bound

We first state a simple general property of the class US.

Lemma 8.1 *Every US-complete set has an AND_2 function.²*

²In fact, a stronger fact is true: every US-complete set has an AND function of unbounded arity.

Proof. It suffices to prove the lemma for any particular US-complete set, for example, USAT. Given two Boolean formulas Φ and Ψ , rename the variables in Ψ so the formulas become over disjoint sets of variables and consider the conjunction $\Phi \wedge \Psi$. As easily seen, the conjunction is in USAT iff both Φ and Ψ are in USAT. ■

In Section 6 with a non-empty set $U \subseteq \{0, 1\}^l$ we associated the number $\text{force}(U)$. Additionally, let us put $\text{force}(\emptyset) = \infty$.

Theorem 8.2 *Let π be a polynomially bounded NP optimization problem. Then $\text{FORCE}_\pi(k) \leq_{\text{dtt}}^{\text{P}} \text{US}$ for each $k \geq 0$.*

Proof. We will assume that π is a minimization problem (the case of maximization problems is quite similar). Suppose that $v(x, y) \leq |x|^c$ for a constant c . Given $1 \leq m \leq |x|^c$, we define

$$\text{sol}^m(x) = \{y \in \text{sol}(x) : v(x, y) = m\}$$

and

$$F_\pi^m(x) = \text{force}(\text{sol}^m(x)).$$

In particular, $F_\pi^m(x) = F_\pi(x)$ if $m = \pi(x)$.

Let k be a fixed integer. Notice that

$$F_\pi(x) \leq k \text{ iff } \bigvee_{m=1}^{|x|^c} (F_\pi^m(x) \leq k \wedge \pi(x) \geq m) \quad (6)$$

(actually, only a disjunction member where $m = \pi(x)$ can be true). The set of pairs (x, m) with $\pi(x) \geq m$ is in coNP and hence in US. Let us now show that the set of (x, m) with $F_\pi^m(x) \leq k$ is dtt-reducible to US.

Recall that $\text{sol}(x) \subseteq \{0, 1\}^{l(x)}$, where $l(x) \leq |x|^d$ for a constant d . Define T to be the set of quadruples (x, m, l, D) such that m and l are positive integers, $D \subseteq [l]$, and there is a unique $y \in \text{sol}^m(x)$ of length l with all 1's in positions from D . It is easy to see that T is in US and

$$F_\pi^m(x) \leq k \text{ iff } \bigvee_{\substack{l, D: l \leq |x|^d \\ D \subseteq [l], |D| \leq k}} (x, m, l, D) \in T.$$

Combining this equivalence with (6), we conclude that $F_\pi(x) \leq k$ iff there are numbers $m \leq |x|^c$ and $l \leq |x|^d$ and a set $D \subseteq [l]$ of size at most k such that

$$(x, m, l, D) \in T \wedge \pi(x) \geq m.$$

By Lemma 8.1, this conjunction is expressible as a proposition about membership of the quadruple (x, m, l, D) in a US-complete set. Thus, the condition $F_\pi(x) \leq k$ is equivalent to a disjunction of less than $|x|^{c+d(k+1)}$ propositions each verifiable in US. ■

Corollary 8.3 *Let $\pi \in \{\chi, \omega, \gamma\}$. Then $\text{FORCE}_\pi(k) \leq_{\text{dtt}}^{\text{P}} \text{US}$ for each $k \geq 0$.*

Remark 8.4 Using (3) and Theorem 8.2, we can easily show that for the vertex cover number τ we also have

$$\text{US} \leq_{\text{m}}^{\text{P}} \text{UO}_\tau = \text{FORCE}_\tau(0) \leq_{\text{m}}^{\text{P}} \text{FORCE}_\tau(k) \leq_{\text{m}}^{\text{P}} \text{FORCE}_\tau(k+1) \leq_{\text{dtt}}^{\text{P}} \text{US}.$$

9 Concluding discussion and open questions

1. We have considered forcing versions of the three most popular graph invariants: the chromatic, the clique, and the domination numbers (F_χ , F_ω , and F_γ respectively). We have shown that the slice decision problems for each of F_χ , F_ω , and F_γ are as hard as US under the many-one reducibility and as easy as US under the dtt-reducibility. The latter upper bound is actually true for the forcing variant of any polynomially bounded NP optimization problem. The lower bound in the case of F_ω and F_γ is provable by using standard reductions on the account of a close connection with the unique optimum problems UO_ω and UO_γ . However, in the case of F_χ we use somewhat more elaborate reductions involving graph products. We point out two simple reasons for the distinction between F_χ and F_ω , F_γ . First, unlike the case of ω and γ , the unique colorability of a graph is apparently inexpressible in terms of F_χ (cf. Lemma 2.7.3). Second, we currently do not know any reductions between F_χ , F_ω , and F_γ as optimization problems that would allow us to relate their complexities (cf. further discussion).

2. We have shown that the slice decision problems $\text{FORCE}_\pi(k)$ for $\pi \in \{\chi, \omega, \gamma\}$ are close to each other in the complexity hierarchy. Furthermore, let $\text{FORCE}_\pi(*) = \{(x, k) : F_\pi(x) \leq k\}$. Hamed Hatami (personal communication) has recently shown that, like $\text{FORCE}_\chi(*)$, the decision problem $\text{FORCE}_\omega(*)$ is Σ_2^P -complete. Consequently, the problems of computing F_χ and F_ω are polynomial-time Turing equivalent. It would be also interesting to compare the complexities of F_χ , F_ω , and F_γ using weaker reducibility concepts for optimization problems (as is well known, the similarity of decision versions does not imply the similarity of the underlying optimization problems; For example, the decision versions of χ , ω , and γ are all NP-complete and parsimoniously equivalent but have pairwise different parametrized complexities and the corresponding optimization problems have pairwise different approximation complexities).

3. Is $\text{FORCE}_\pi(k)$ NP-hard under $\leq_m^P$ -reductions for any π under consideration and constant k ? It should be noted that the affirmative answer would settle a long-standing open problem if C=P contains NP in the affirmative.

4. Let UCOL be the set of all uniquely colorable graphs (with no restriction on the chromatic number). Is it true that $\text{UCOL} \leq_m^P \text{FORCE}_\chi(2)$? It is not hard to show that UCOL is US-hard.

Acknowledgement This work was initiated when the authors met at the Department of Information Systems at Vienna University of Technology. They are indebted to Georg Gottlob for his hospitality.

In Memoriam The journal version of this paper is prepared already without Frank Harary. He passed away a month before the conference presentation at STACS'05 [24]. Frank's enthusiasm and inspiration were a fundamental of this work.

W.S., O.V.

References

- [1] P. Adams, M. Mahdian, E.S. Mahmoodian. On the forced matching numbers of bipartite graphs. *Discrete Mathematics* 281:1–12 (2004).
- [2] A.V. Aho, J.E. Hopcroft, J.D. Ullman. *The design and analysis of computer algorithms*. Addison-Wesley (1976).
- [3] P. Afshani, H. Hatami, E.S. Mahmoodian. On the spectrum of the forced matching number of graphs. *Australas. J. Combin.* 30:147–160 (2004).
- [4] A. Blass, Y. Gurevich. On the unique satisfiability problem. *Information and Control* 55:80–88 (1982).
- [5] R. Chang, J. Kadin, P. Rohatgi. On unique satisfiability and the threshold behaviour of randomized reductions. *Journal of Computer and System Sciences* 50(3):359–373 (1995).
- [6] G. Chartrand, H. Gavlas, R.C. Vandell, F. Harary. The forcing domination number of a graph. *J. Comb. Math. Comb. Comput.* 25:161–174 (1997).
- [7] G. Chartrand, F. Harary, M. Schultz, C.E. Wall. Forced orientation numbers of a graph. *Congr. Numer.* 100:183–191 (1994).
- [8] G. Chartrand, P. Zhang. The forcing geodetic number of a graph. *Discuss. Math. Graph Theory* 19:45–58 (1999).
- [9] G. Chartrand, P. Zhang. The forcing hull number of a graph. *J. Comb. Math. Comb. Comput.* 38:81–94 (2001).
- [10] G. Chartrand, P. Zhang. The forcing dimension of a graph. *Mathematica Bohemica* 126(4):711–720 (2001).
- [11] C.J. Colbourn, M.J. Colbourn, D.R. Stinson. The computational complexity of recognizing critical sets. *Graph theory, Proc. 1-st Southeast Asian Colloq.* Singapore 1983, Lect. Notes Math., Vol. 1073, 248–253 (1984).
- [12] P. Crescenzi. A short guide to approximation preserving reductions. In: *Proc. of the 12-th Ann. Conference on Computational Complexity*, 262–272 (1997).
- [13] P. Crescenzi, C. Fiorini, R. Silvestri. A note on approximation of the MAX CLIQUE problem. *Information Processing Letters* 40:1–5 (1991).
- [14] D. Curran, G.H.J. van Rees. Critical sets in latin squares. In: *Proc. of the 8-th Manitoba conference on numerical mathematics and computing, Winnipeg 1978. Congressus Numerantium* 22:165–168 (1979).
- [15] D.P. Dailey. Uniqueness of colorability and colorability of planar 4-regular graphs are NP-complete. *Discrete Math.* 30, 289–293 (1980).
- [16] T. Easton, R. Gary Parker. On completing latin squares. *Discrete Appl. Math.* 113(2-3):167–181 (2001).
- [17] M.R. Garey, D.S. Johnson, L.J. Stockmeyer. Some simplified NP-complete graph problems. *Theor. Comp. Sci.* 1:237–267 (1976).
- [18] K. Gray. On the minimum number of blocks defining a design. *Bull. Aust. Math. Soc.* 41(1):97–112 (1990).
- [19] D.L. Greenwell, L. Lovász. Applications of product colouring. *Acta Math. Acad. Sci. Hung.* 25:335–340 (1974).

- [20] H. Hajiabolhassan, M.L. Mehrabadi, R. Tusserkani, M. Zaker. A characterization of uniquely vertex colorable graphs using minimal defining sets. *Discrete Math.* 199:233–236 (1999).
- [21] F. Harary. Three new directions in graph theory. In: *Proceedings of the First Estonian Conference on Graphs and Applications* (Tartu-Kääriku, 1991). Tartu: Tartu Univ. 15–19 (1993).
- [22] F. Harary, S.T. Hedetniemi, R.W. Robinson. Uniquely colorable graphs. *J. Combin. Theory* 6:264–270 (1969); 9:221 (1970).
- [23] F. Harary, D. Klein, T. Živković. Graphical properties of polyhexes: Perfect matching vector and forcing. *J. Math. Chemistry* 6:295–306 (1991).
- [24] F. Harary, W. Slany, O. Verbitsky. On the computational complexity of the forcing chromatic number. In: *Proc. of the 22nd Annual Symposium on Theoretical Aspects of Computer Science*. V. Diekert, B. Durand Eds. Lecture Notes in Computer Science, Vol. 3404. Berlin: Springer-Verlag. 182–193 (2005).
- [25] H. Hatami, H. Maserrat. On the computational complexity of defining sets. *Discrete Applied Mathematics*, to appear.
- [26] L. Hemachandra. Structure of complexity classes: separations, collapses, and completeness. In: *Proc. of the 13-th MFCS conf.* Lecture Notes in Computer Science, Vol. 324, 59–73 (1988).
- [27] L.A. Hemaspaandra, M. Ogihara. *The complexity theory companion*. Berlin: Springer (2002).
- [28] P. Horák, R.E.L. Aldred, H. Fleischner. Completing Latin squares: Critical sets. *Journal of Combinatorial Designs* 10(6):419–432 (2002).
- [29] W. Imrich, S. Klavzar. *Product graphs. Structure and recognition*. Wiley-Interscience Series in Discrete Mathematics and Optimization. Wiley, Chichester (2000).
- [30] V. Kann. *On the approximability of NP-complete optimization problems*. Ph.D. thesis. Royal Institute of Technology, Stockholm, 1992.
- [31] J. Köbler, U. Schöning, and J. Torán. *The Graph Isomorphism problem: its structural complexity*. Birkhäuser, 1993.
- [32] C. Lund, M. Yannakakis. On the hardness of approximating minimization problems. *J. of the ACM* 41(5):960–981 (1994).
- [33] M. Mahdian, E.S. Mahmoodian, R. Naserasr, F. Harary. On defining sets of vertex colorings of the Cartesian product of a cycle with a complete graph. In: *Combinatorics, Graph Theory, and Algorithms*, Y. Alavi, D.R. Lick, and A. Schwenk Eds. 461–467. New Issues Press, Kalamazoo (1999).
- [34] E.S. Mahmoodian, R. Naserasr, M. Zaker. Defining sets in vertex colorings of graphs and Latin rectangles. *Discrete Math.* 167-168:451–460 (1997).
- [35] J. Nelder. Critical sets in latin squares. *CSIRO Division of Math. and Stats, Newsletter* 38:4 (1977).
- [36] C. Papadimitriou. On the complexity of unique solutions. *J. of the ACM* 31(2):392–400 (1984).
- [37] C. Papadimitriou. *Computational complexity*. Addison-Wesley, 1994.

- [38] D. Ranjan, S. Chari, and P. Rohatgi. Improving known solutions is hard. *Computational Complexity* 3(2):168–185 (1993).
- [39] A.L. Selman. Promise problems and complexity classes. *Information and Computation* 78:87–98 (1988).
- [40] L.J. Stockmeyer. Planar 3-colorability is NP-complete. *SIGACT News* 5(3):19–25 (1973).
- [41] T. Thierauf. *The computational complexity of equivalence and isomorphism problems*. Lecture Notes in Computer Science, Vol. 1852. Springer (2000).
- [42] L.G. Valiant, V.V. Vazirani. NP is as easy as detecting unique solutions. *Theor. Comp. Sci.* 47(1):287–300 (1986).
- [43] D. Zuckerman. On unapproximable versions of NP-complete problems. *SIAM J. Comput.* 25(6):1293–1304 (1996).